

Cimarron Basin High Priority Issue


Bioenergy and Water

January 2009

onomic practices and crop genetics have led to higher corn yields. In southwest Kansas, while there was a 16% increase in irrigated corn acres from 1993 to 2003, there was a 52% increase in irrigated corn production.⁽¹⁰⁾ According to the Kansas Department of Agriculture, Stevens County ranks first in all Kansas counties for irrigated corn acres.

In 2006, approximately 16% of Kansas corn and sorghum [crops](#) were used for ethanol production, up 13% from 2000. Corn production in Kansas may be slowing down. According to NASS, producers intended to plant eight percent fewer corn acres in 2008, as a result of multiple factors including crop rotation considerations and high input costs. In 2008, Kansas was expected to plant their largest soybean crops in history.⁽⁵⁾

Water Quality



Wastewater from ethanol plants is regulated by the Kansas Department of Health and Environment (KDHE), which administers both the federal National Pollution Discharge Elimination System (NPDES) permits and Kansas Water Pollution Control permits. In most instances, KDHE issues the state-level permit, which requires ethanol plants to use the wastewater for beneficial land applications rather than simply discharging into streams and rivers.

A rise in the number of corn acres may also impact the basin's water quality through increased fertilizer application and soil erosion. Corn has the greatest application rates of both fertilizer and pesticides per acre, higher

than for soybeans and mixed-species grassland biomass. The switch from other crops or noncrop plants to corn may lead to higher application rates of highly soluble nitrogen. Harvested row crops, such as corn, have a higher potential for soil erosion than grasses or perennial crops. The potential water quality impact of an increased demand for corn may be mitigated through Best Management Practices (BMPs), especially those addressing soil erosion and herbicide applications.

Biodiesel

Biodiesel is produced using oils extracted from crops, animal fat or waste vegetable oil using a chemical process called transesterification. Most U.S. biodiesel is produced from soybean oil, although other vegetable oils such as canola, corn, cottonseed, flax seed, sunflower, or peanut oil can be used. As of December 2008, no biodiesel facilities are permitted for or located in the Cimarron basin.

Biodiesel production uses roughly three gallons of water per gallon, about a gallon of which is consumptive use. Wastewater from biodiesel plants, which may contain high amounts of oxygen, grease and oils, is regulated by the KDHE.

Cellulosic Ethanol

Cellulosic ethanol uses lignocellulose, the main structural material in any plant, as a feedstock. Cellulosic feedstocks require an extra step to break down the lignocellulose into fermentable starch, thus increasing production costs. The bulkier cellulosic feedstocks are also more costly to harvest, transport and store. Processing of cellulosic

materials would require more water than corn, as the feedstock is dry. Research on cellulosic feedstocks, such as switchgrass, wood chips, and corn stover is ongoing. The U.S. Department of Energy has set 2012 as a target to achieve technological advances to make cellulosic ethanol cost competitive with corn ethanol. In 2007, Abengoa Bioenergy, a Spanish energy company, announced that Hugoton, Kansas would be the site of the state's first cellulosic ethanol plant. In conjunction with cellulosic ethanol research, some researchers are investigating the use of perennial polyculture crop systems for cellulosic feedstocks.

Production of cellulosic ethanol may have greater positive environmental impacts than grain based ethanol

Cimarron Basin High Priority Issue

Bioenergy and Water

January 2009

Page 3

such as reduced greenhouse gas emissions, decreased fertilizer application and less reliance on water intensive crops.

Corn Research and Varieties

Breeding of corn hybrids that maximize yield for ethanol production while reducing additional strains on water supplies has been a focus of much research by universities and corn breeding companies. Drought tolerant hybrids, specifically transgenic drought resistant corn, are especially important in areas of western Kansas where rainfall averages fewer than 16 inches per year. In addition to drought tolerant varieties, industries are identifying corn varieties that produce higher yield and more ethanol per acre. High total fermentable ethanol corn hybrids provide higher levels of fermentable starch, the sum of all starches and simple sugars that ferment during the typical dry grind process.^(4,7)



Recommended Actions

1. Coordinate, where applicable, the development, implementation, and public input process between the *Kansas Water Plan* and Kansas energy policy.
2. Maintain regulatory oversight by state and local government on the siting of ethanol and biodiesel plants, with special emphasis on educating applicants on the water supply and availability.
3. Look for water recycling opportunities within the bio-fuel facilities.
4. Promote research for less water dependent corn varieties and improved irrigation scheduling that maintains or increases crop yield without increasing water use, and encourage more use of grain sorghum, sweet sorghum and other lower water use crops.
5. Promote research and pilot projects for viable, commercial cellulosic ethanol production and other bio-fuels less dependent on water intensive crop production.
6. Increase corn water use efficiency (amount of grain produced per inch of water) through research and extension efforts. Educational emphasis should be placed on utilization of irrigation scheduling tools such as KanSched and the Mobile Irrigation Lab.
7. Evaluate the biofuel facility watershed and watersheds of input crops, and identify potentially environmentally sensitive areas and target programs to mitigate environmental impacts, such as stream buffers, grass filters, BMPs, etc.
8. Provide education and/or incentives for marginal lands that have expiring, un-renewable Conservation Reserve Program contracts to stay in a conservation planting, with special consideration to acres that could return to irrigation.

Cimarron Basin High Priority Issue

Bioenergy and Water

January 2009

Page 4

Resources

1. Kansas Corn Growers Association. April 2008. *Kansas Corn and Water Quality*. <http://www.ksgains.com/corn/water.html>
2. Kansas Energy Council. December 2007. *Kansas Energy Report 2008*.
3. Kansas Energy Council. January 2007. *Kansas Energy Report 2007*. *Kansas Energy Chart Book*.
4. Monsanto Industry Webpage, accessed on May 6, 2008. http://monsanto.com/products/seeds_traits/corn.asp
5. National Agricultural Statistics Service (NASS), Agricultural Statistics Board, U.S. Department of Agriculture. March 2008. *Prospective Plantings*.
6. National Agricultural Statistics Service (NASS), Agricultural Statistics Board, U.S. Department of Agriculture. March 2008. *Crops and Plants National Statistics*.
7. Pioneer Industry Webpage, accessed on May 6, 2008. www.pioneer.com
8. Science Daily. Retrieved April 4, 2008. <http://www.sciencedaily.com/releases/2007/10/071010120538.htm>
9. U.S. Department of Energy. Retrieved 2008. <http://www.doe.gov/energysources/bioenergy.htm>
10. White, Jere, Executive Director, Kansas Corn Growers, August 13, 2008, written communication.

